



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

III. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Suppose $ABCD$ the required trapezoid, AB being the shorter (upper) base, $BD=x$, one diagonal, $AC=BF=y$, $BG=\text{altitude}=h$. Produce DC to F so that $DF=DG+GF=DC+CF=a$, $x+y=c$, $\theta=\text{given angle}$. $\sqrt{(x^2-h^2)}+\sqrt{(y^2-h^2)}=a$. $4x^2y^2=a^4+(x^2+y^2)^2+4a^2h^2-2a^2(x^2+y^2)$. But $x^2+y^2=c^2-2xy$. $\therefore 4(c^2-a^2)xy=4a^2h^2+(c^2-a^2)^2$. Now $xy\sin\theta=ah$.

$$\therefore 4a^2h^2-4ah\left(\frac{c^2-a^2}{\sin\theta}\right)=-(c^2-a^2)^2.$$

$$\therefore h=\frac{1}{2a}(c^2-a^2)\tan\frac{1}{2}\theta \text{ or } h=\frac{1}{2}(c^2-a^2)\cot\frac{1}{2}\theta.$$

$$\therefore xy=\frac{c^2-a^2}{4\cos^2\frac{1}{2}\theta} \text{ or } \frac{c^2-a^2}{\sin^2\frac{1}{2}\theta} \text{ and } x+y=c.$$

$$\therefore x=\frac{1}{2}c\pm\frac{\sqrt{(a^2-c^2\cos^2\frac{1}{2}\theta)}}{2\sin\frac{1}{2}\theta} \text{ or } \frac{1}{2}c\pm\frac{\sqrt{(a^2-c^2\sin^2\frac{1}{2}\theta)}}{2\cos\frac{1}{2}\theta};$$

$$y=\frac{1}{2}c\mp\frac{\sqrt{(a^2-c^2\cos^2\frac{1}{2}\theta)}}{2\sin\frac{1}{2}\theta} \text{ or } \frac{1}{2}c\mp\frac{\sqrt{(a^2-c^2\sin^2\frac{1}{2}\theta)}}{2\cos\frac{1}{2}\theta}.$$

Hence lay off $DF=a$, and draw AE parallel to DF at a distance h from it. With D as center and $DB=x$ or y draw DB ; then $DF=y$ or x . Draw any line parallel to BF as $AC, A'C', A''C''$, and join B, D to C, C', C'', A, A', A'' , respectively. Then any one of the many trapezoids thus formed fulfill the required conditions, as is evident by drawing a figure.

IV. Solution by J. J. KEYES, Nashville, Tenn.

Construct the triangle EBF having $BE=\text{sum of diagonals}$, $BF=\text{sum of bases}$, and angle $E=\frac{1}{2}$ given angle. At F construct the angle $EFD=\text{angle } E$, FD meeting BE in D . Through D draw DM parallel to FB . Take any point C on BF , draw CA parallel to FD meeting DM in A . Join AB, DC . $ABCD$ is the required trapezoid. In proof, $AO=CF$. $\therefore AD+BC=BC+BD=\text{sum of bases}$. $AC=DF=DE$. $\therefore AC+BD=BD+DE=\text{sum of diagonals}$. The angle between BD and $AC=\text{angle } BDF=?$ angle $E=\text{given angle}$.

Also solved by Elmer Schuyler.

CALCULUS.

Problem number 181 was also solved by S. A. Corey and L. C. Walker.

183. Proposed by W. J. GREENSTREET, A. M., Stroud, England.

Evaluate $\int_0^\infty \frac{\sin 2nxdx}{(a^2+x^2)\sin x}$.

Solution by G. B. M. ZERR, A. M., Ph. D.

Let $u = \int_0^\infty \frac{\sin 2nx \, dx}{(a^2 + x^2) \sin x}$, then

$$\frac{du}{dn} = \int_0^\infty \frac{2x \cos nx \, dx}{(a^2 + x^2) \sin x}, \quad \frac{d^2 u}{dn^2} = - \int_0^\infty \frac{4x^2 \sin 2nx \, dx}{(a^2 + x^2) \sin x} = 4a^2 u - 4 \int_0^\infty \frac{\sin 2nx \, dx}{\sin x}.$$

$$\text{Let } v = 8 \int_0^\infty \frac{\sin nx \cos nx \, dx}{\sin x} = 8 \int_0^\infty \frac{e^{nx\sqrt{-1}} - e^{-nx\sqrt{-1}}}{e^{x\sqrt{-1}} - e^{-x\sqrt{-1}}} \cos nx \, dx.$$

Let $x\sqrt{-1} = \pi y$, $dx = -\pi\sqrt{-1} dy$;

$$\begin{aligned} \therefore v &= -8\pi\sqrt{-1} \int_0^\infty \frac{e^{\pi ny} - e^{-\pi ny}}{e^{\pi y} - e^{-\pi y}} \cos(\pi ny\sqrt{-1}) dy \\ &= -8\pi\sqrt{-1} \left(\frac{\sin \pi n}{e^{\pi n\sqrt{-1}} + 2\cos \pi n + e^{-\pi n\sqrt{-1}}} \right) = -2\pi\sqrt{-1} \tan \pi n. \end{aligned}$$

$$\therefore v = 0, \text{ since } n = \text{any integer.} \quad \therefore \frac{d^2 u}{dn^2} - 4a^2 u = 0.$$

$u = Ae^{2an} + Be^{-2an}$. When $n=0$, $u=0$, and $B=-A$. $\therefore u = A(e^{2an} - e^{-2an})$.

When $n=1$, $u = \frac{\pi e^{-a}}{a}$ and $A = \frac{\pi e^{-a}}{a(e^{2a} - e^{-2a})}$.

$$\therefore u = \frac{\pi e^{-a}}{a} \cdot \frac{e^{2an} - e^{-2an}}{e^{2a} - e^{-2a}} = \frac{\pi e^{-a}}{a} \cdot \frac{\sinh 2an}{\sinh 2a}.$$

184. Proposed by W. J. GREENSTREET, A. M., Stroud, England.

If $u = f(x, y)$; $\xi = e^x y$; $\eta = e^x$; show that

$$\frac{d^2 u}{dx^2} - y^2 \frac{d^2 u}{dy^2} - y \frac{du}{dy} = 4\xi \eta \frac{d^2 u}{d\xi d\eta}.$$

Solution by G. W. GREENWOOD, M. A. (Oxon), Lebanon, Ill.

We have $x = \frac{1}{2} \log \xi + \frac{1}{2} \log \eta$, $y = \sqrt{\xi/\eta}$, and therefore

$$\frac{\partial}{\partial \xi} = \frac{\partial x}{\partial \xi} \cdot \frac{\partial}{\partial x} + \frac{\partial y}{\partial \xi} \cdot \frac{\partial}{\partial y} = \frac{1}{2} \left(\frac{1}{e^x y} \frac{\partial}{\partial x} + \frac{1}{e^x} \frac{\partial}{\partial y} \right), \text{ whence } 2\xi \frac{\partial}{\partial \xi} = \frac{\partial}{\partial x} + y \frac{\partial}{\partial y}.$$

In a similar way we get

$$2\eta \frac{\partial}{\partial \eta} = \frac{\partial}{\partial x} = y \frac{\partial}{\partial y}.$$